

Properties of Stellar Clusters around High-Mass Young Stars[★]

F. Faustini¹, S. Molinari¹, L. Testi^{2,3} and J. Brand⁴

¹ Istituto di Fisica dello Spazio Interplanetario - INAF, via Fosso del Cavaliere 100, I-00133, Rome (e-mail: faustini, molinari@ifsi-roma.inaf.it)

² European Southern Observatory, Karl Schwarzschild str. 2, Garching bei Muenchen, Germany (e-mail: ltesti@eso.org)

³ Osservatorio Astronomico di Arcetri - INAF, via Enrico Fermi 5, I-50125 Firenze

⁴ Istituto di Radioastronomia - INAF, Via Gobetti 101, Bologna (e-mail: brand@ira.inaf.it)

Received ; accepted

ABSTRACT

Context. Twenty-six high-luminosity IRAS sources believed to be collection of stars in the early phases of high-mass star formation have been observed in the Near-IR (J, H, K_s) to characterize the clustering properties of their young stellar population and compare them with those of more evolved objects (e.g., Herbig Ae/Be stars) of comparable mass. All the observed sources possess strong continuum and/or line emission in the millimeter, being therefore associated with gas and dust envelopes. Nine sources have far-IR colors characteristic of UCHII regions while the other 17 are likely being experiencing an evolutionary phase that precedes the Hot-Cores, as suggested by a variety of evidence collected in the past decade.

Aims. To gain insight into the initial conditions of star formation in these clusters (Initial Mass Function [IMF], Star Formation History [SFH]), and to deduce mean values for cluster ages.

Methods. For each cluster we carry out aperture photometry. We derive stellar density profiles, color-color and color-magnitude diagrams, and color (HKCF) and luminosity (KLF) functions. These two functions are compared with simulated KLFs and HKCFs from a model that generates populations of synthetic clusters starting from assumptions on the IMF, the SFH, and the Pre-MS evolution, and using the average properties of the observed clusters as boundary conditions (bolometric luminosity, dust distribution, infrared excess, extinction).

Results. Twenty-two sources show evidence of clustering with a stellar richness indicator that varies from a few up to several tens of objects, and a median cluster radius of 0.7 pc. A considerable number of cluster members present an infrared excess characteristic of young Pre-Main-Sequence objects. For a subset of 9 detected clusters, we could perform a statistically significant comparison of the observed KLFs with those resulting from synthetic cluster models; for these clusters we find that the median stellar age ranges between $2.5 \cdot 10^5$ and $5 \cdot 10^6$ years, with evidence of an age spread of the same entity within each cluster. We also find evidence that older clusters tend to be smaller in size, in line with the fact that our clusters are on average larger than those around relatively older Herbig Ae/Be stars. Our models allow us to explore the relationship of the mass of the most massive star in the cluster with both the clusters richness and their total stellar mass. Although such relationships are predicted by several classes of cluster formation models, their detailed analysis suggests that our modeled clusters may not be consistent with them resulting from random sampling of the IMF.

Conclusions. Our results are consistent with a star formation which takes place continuously over a period of time which is longer than a typical crossing time.

Key words. Stars: formation – Stars: imaging – Stars: luminosity function, mass function – Stars: pre-main sequence – Infrared: stars

1. Introduction

The last few decades have been characterized by a large effort to improve our understanding of how stars form both from a theoretical and from an observational point of view. As a result, today we have reached a good understanding of how isolated low-mass stars form (Klein et al. 2006). The widely accepted scenario is that low mass stars form through the gravitational collapse of a prestellar core followed at later stages by disk accretion.

Extending this theory to high-mass stars is not trivial. High mass (proto-)stars reach the Zero Age Main Sequence while still accreting. When the central protostar reaches about $10 M_{\odot}$ hydrogen fusion ignites in the core and the star's radiation pressure and wind should prevent further accretion. This obviously is a paradox given that more massive stars do form. Several theo-

ries have been put forward to solve this dilemma (Zinnecker & Yorke 2007): accretion rates as high as three orders of magnitude larger than in the case of low-mass stars (Cesaroni 2005), and non-spherical accretion geometries (Nakano 1989, Yorke 2002, Keto 2003), or coalescence in dense (proto-)stellar clusters (Bonnell et al. 1998).

All these theories have predictions that, in principle, could be tested observationally. In the last decade a large effort was made in trying to detect massive accretion disks (Cesaroni et al. 2006), powerful outflows (Beuther et al. 2002, Cesaroni et al. 2005) and dense protostellar clusters (Testi et al. 1999, de Wit et al. 2005), all of these phenomena are predicted by one or the other formation theory. None of these efforts have provided conclusive arguments in favour or against any of the theories.

In this paper we explore the properties of embedded clusters associated with high-mass protostellar candidates. Our sample was selected from a larger sample of candidate high-mass protostars selected and analyzed, in the past decade, by Molinari et

Send offprint requests to: faustini@ifsi-roma.inaf.it

[★] Based on observations obtained at the Palomar Observatory and at the ESO La Silla Observatory (Chile), programme 65.I-0310(A)

al. (1996, 1998, 2000, 2002) and Brand et al. (2001). In Sect. 2 we present the observations and data analysis (source extraction, photometry), in Sect. 3 we discuss data elaboration and interpretation. In Sect. 4 we present our Synthetic Cluster Generation model and the method of comparison between synthetic and observed clusters and the results of using this technique. Finally in Sect. 5 we compare our objects with more evolved ones and present our conclusions.

2. Observations and Data Analysis

Program fields are listed in Table 1 and were imaged in J, H, and K_s bands. A total of 15 fields were observed in three nights in November 1998 at the Palomar 60-inch telescope equipped with a 256×256 NICMOS-3 array, with a pixel scale of $0''.62/\text{pix}$ and total FOV of $2'.6 \times 2'.6$. The remaining 11 fields were observed in 3 nights in August 2000 at the ESO-NTT using the 1024×1024 SOFI camera, with a pixel scale of $0''.29/\text{pix}$ and a total FOV of $4'.9 \times 4'.9$. Standard dithering techniques were used to minimize impact of bad pixels and optimize flat-fielding, allowing to achieve for each field a total of 5min integration time per band (in the central portion of the observed field) for a covered area of $3'.5 \times 3'.5$ for Palomar observations, and 20min (10min for the band K_s) at NTT with a total covered area of $6'.5 \times 6'.5$. Suitable calibration sources from the list of Hunt et al. (1988) were observed regularly during the observations to track atmospheric variations for different airmasses. Standard stars and target fields were observed at airmasses no greater than 1.7 at NTT, and 1.3 at Palomar; we determined average zero-point magnitudes for each night and used them to calibrate our photometry. For each field the images in the three bands were registered and astrometric solutions were determined using a few bright optically visible sources.

The K_s images for all observed fields, with superimposed submillimeter continuum emission distribution when available (Molinari et al. 2008a) are presented in Appendix A and available online at <http://galatea.ifs-roma.inaf.it/faustini/K-images>.

2.1. Point Source Extraction and Photometry

The extraction and photometry of point sources for all images was carried using the IRAF package. The r.m.s of the background signal and the FWHM of point sources were measured throughout the images to characterize the image noise and PSF properties; these parameters were fed to the DAOFIND task for source extraction, where a detection threshold of 3σ was used for all images. Sources with saturated pixels were excluded from the analysis; the linearity of the system response was checked *a posteriori* comparing, both for the Palomar and the NTT data, the magnitudes obtained to those from 2MASS using a few stars with magnitudes reaching up to the maximum values found in our photometry files; the relations between the 2MASS magnitudes and ours in the three bands were found to be linear over the entire magnitude range of the detected sources. There were clearly brighter objects in the various fields, but their peaks were already flagged as saturated and were excluded from the detection process.

The photometry of sources is critical in very dense stellar fields like the inner Galactic Plane, where all of our target fields lie, and where the crowding is such that more than one source can enter any plausible aperture that can be chosen, or any annulus used for background estimation. This problem is of course magnified in the clustered environments that are detected in sites of massive star formation (see §3.1 below).

Table 1. Journal of Observations

Source Mol ^a	IRAS Name	α (J2000)	δ (J2000)	Tel.
3	00420+5530	00:44:57.6	+55:46:52	Pal
8	05137+3919	05:17:13.3	+39:22:14	Pal
9	05168+3634	05:20:16.2	+36:37:21	Pal
11	05345+3157	05:37:47.8	+31:59:24	Pal
12	05373+2349	05:40:24.4	+23:50:54	NTT
15	06056+2131	06:08:41.0	+21:31:01	Pal
28	06584-0852	07:00:51.0	-08:56:29	Pal
30	17450-1742	17:48:09.3	-27:43:21	NTT
38	18024-2119	18:06:18.0	-21:42:00	NTT
45	18144-1723	18:17:24.2	-17:22:13	NTT
50	18162-1612	18:19:07.5	-16:11:21	NTT
59	18278-1009	18:30:35.2	-10:07:12	Pal
75	18511+0146	18:53:38.1	+01:50:27	Pal
82	18565+0349	18:59:03.4	+03:53:22	NTT
84	18567+0700	18:59:13.6	+07:04:47	NTT
98	19092+0841	19:11:37.4	+08:46:30.0	NTT
99	19094+0944	19:11:52.0	+09:49:46	Pal
103	19213+1723	19:23:37.0	+17:28:59	NTT
109	19374+2352	19:39:33.2	+23:59:55	NTT ^b
110	19388+2357	19:40:59.4	+24:04:39	NTT ^b
136	21307+5049	21:32:31.5	+51:02:22.0	Pal
139	21519+5613	21:53:38.8	+56:27:53.0	Pal
143	22172+5549	22:19:09.0	+56:04:45.0	Pal
148	22305+5803	22:32:24.3	+58:18:58.2	Pal
151	22506+5944	22:52:38.6	+60:00:56.0	Pal
160	23385+6053	23:40:53.3	+61:10:19.1	Pal

^a Source running number from Molinari et al. 1996.

^b Imaged only in K_s .

The first approach we tried to follow is the PSF-fitting photometry which should be less prone to these problems. We chose a sub-sample of test fields with different levels of stellar crowding. An important aspect in this procedure is the modelling of the PSF; we made several trials selecting a variable number of point-like sources (from 3 to 30) of different brightness levels and different position in the field. We find that the resulting PSF model is not particularly sensitive to the choice of numbers and/or brightness of the stars; on the other hand the results are quite different depending on the mean stellar density of the field. The photometry was carried out using the ALLSTAR task, particularly suited for crowded fields. However, we also tested the other two tasks (PEAK and NSTAR) and they produce comparable results for most of the sources. We note, however, that especially in the most crowded areas the subtraction of the PSF-fitted sources from the image introduces two spurious effects: an unacceptably high level of residuals with brightness levels well above the detection threshold used and a great number of negative holes which show that the psf-fit includes a portion of background in the estimation of the source flux, therefore overestimating its value. This is due both to the limited accuracy of the PSF model that can be built in very crowded fields, where faint neighboring stars can enter in the area where the PSF model is estimated, and to the presence of a significant and variable background which is quite common and expected in the Galactic Plane. A similar conclusion was reached by Hillenbrand & Carpenter (2000) in their study of the inner Orion Nebula Cluster.

The second approach we followed is standard aperture photometry. The choice of the radii for the aperture and for the background annuli is of course extremely important. The optimum aperture should not be too large to include nearby sources and not too small to significantly cut the PSF and severely under-

estimate the flux. We did several attempts on one of the most crowded fields (Mol30, observed at NTT) with three different aperture radii equal to the PSF FWHM (typically $0''.7$ at NTT and $1''.4$ at Palomar in K_s), twice and thrice this value. For each photometry run we analyze the sources' flux distribution and, as expected, the median flux is found to increase with increasing aperture radius. Increasing the aperture from one to two PSF FWHMs raises the median source flux by an amount compatible with the inclusion of the first ring of an Airy diffraction pattern; instead, an aperture radius equal to three times the PSF FWHM produces a flux increase much too large compared to the additional fraction of the Airy profile entering the aperture, therefore being caused by the inclusion of nearby sources. We adopted an aperture radius equal to the PSF FWHM to minimize neighbour contamination, and then applied an aperture correction factor for the fraction of the PSF cut out of the aperture; this was estimated via multi-aperture photometry (starting from a 1 FWHM size) on relatively isolated stars in the target fields.

A further effect to be corrected for, also given the crowding of our fields, is the possible contamination arising from the tails of the brightness profiles of neighbouring stars. To quantify this contamination we created a grid of simulations with two symmetric Gaussians with a wide variety of peak contrast and at different reciprocal distances. We computed the fraction of the Gaussian profile of the neighbouring source that falls into the photometry aperture centered on the main source, and hence generated a matrix of photometry corrections for different source distances and peak contrasts. At this point we run through the magnitude file produced by the aperture photometry task and for each source we apply a magnitude correction depending on the presence, distance and contrast ratios with other neighbouring stars.

In spite of the various issues discussed above, the photometry obtained with the two methods are in a good agreement with each other, except at faint magnitudes. For these faint objects we always find that the PSF photometry tends to produce lower magnitudes (hence stronger sources) than the aperture photometry; this effect is easily understood given our finding (see above) that the subtraction of PSF-fitted sources always leaves negative holes in the residual image, and this effect is much more important for faint stars. We thus decided to adopt the magnitudes determined from aperture photometry.

For each target field we estimated the limiting magnitude (LM) using artificial star experiments. The fields were populated using the IRAF task ADDSTAR with 400 fake stars with magnitudes distributed in bins of 0.25 mag between values of 15 and 21; the percentage of recovered stars as a function of magnitude gives an estimate of the completeness level of our photometry. The star recovery percentage does not decrease monotonically with increasing magnitude because fake stars can be placed also very close to bright real stars and then go undetected by the finding algorithm. However, we find that the limit of 85-90% recovery fraction is reached on average around $J=18.7$, $H=17.7$ and $K_s=17.4$ for NTT images, and $J=18.0$, $H=17.3$ and $K_s=16.6$ for Palomar images. We find that the typical photometric uncertainty is below 0.1 mag close to the limiting magnitude.

To verify the integrity of our photometry we compared our magnitudes with those extracted from 2MASS point source catalog for all the fields of our sample. Considering the different spatial resolutions between 2MASS and the telescopes used for our observations, this comparison was limited to those 2MASS point-like sources associated with a single source in the Palomar or NTT images. The median differences with respect to 2MASS for the various fields are of the order of -0.1 , -0.2 and -0.3 mag

for J, H, and K_s bands, respectively. Within each field, the scatter around these median values is ~ 0.1 mag in all three bands, confirming the internal consistency of our photometry. Noticeable departures (~ 0.5 mag) of the median difference with 2MASS from the above values are observed for the field of source Mol11 (Palomar), and for sources Mol103, Mol109 and Mol110 (NTT). However, the latter sources were observed on the same night, which our log registered as not good due to sky variations which are not tracked by night-averaged zero-points. We stress again, however, that these are systematic differences with respect to 2MASS in this limited number of cases; the r.m.s. scatter around these median differences are ~ 0.1 mag in all bands and this should give confidence that the internal consistency of the photometry in each field is preserved. We then decided to rescale our photometry to the 2MASS photometric system to remove these systematic effects. The (J-H) and (H-K) color differences between 2MASS and our photometry are not correlated with the magnitude, so that no magnitude-dependent color effect is introduced in this rescaling.

3. Results

3.1. Cluster Identification

The identification of a cluster results from the analysis of stellar density in the field. Since our target fields are sites of massive star formation associated with local peaks of dust column densities and hence of visual extinction, the K_s images are clearly more suited for this type of analysis.

Stellar density maps were built for each field counting stars in a running boxcar of size equal to $20''$. The box size was determined empirically to enhance the statistical significance of local stellar density peaks and to maximize the ability to detect the clusters. Larger boxes tend to smear the cluster into the background stellar density field decreasing the statistical significance of the peak, which may lead to non-detection of a clearly evident cluster, particularly in the rich inner Galaxy fields (this happens, e.g., for source Mol103, see Fig. A.18 in the Appendix). Smaller boxes produce noisy density maps where the number of sources in each bin starts to be comparable to the fluctuations of the background density field due either to intrinsic variations of the field star density or to variable extinction from diffuse foreground ISM in the Galactic Plane (where all of our sources are located). For most of our objects in the outer Galaxy this analysis is just used to locate the position of the peak stellar density, since the clusters are obvious already from the visual inspection (Mol3 to Mol28, and Mol143 to Mol151, see Appendix). For the rest of the fields the density maps are used to ascertain the presence of a cluster; especially toward the inner Galaxy the density maps tend to show more than one peak at comparable levels. It is important to remember, however, that this is a search for stellar clusters toward regions where indications of active star formation are already available, and this information can be used. In particular, the coincidence of these peaks with cold dust clumps as traced by intense submillimeter and millimeter emission (Beltrán et al. 2006, Molinari et al. 2008a) is critical to consider the density peak as a real feature associated with the star formation region. Casual association is ruled out by the high number of positive associations (see Table 2).

As a further confirmation step for the positive detection of a cluster we build radial stellar density profiles where stars are counted in annuli of increasing internal radius and constant width and then ratioed to the area of the annuli (Testi, Palla & Natta 1998); uncertainties are assigned assuming Poisson statis-




Fig. 1. Stellar density (in stars/pc²), for Mol28, as a function of the radial distance (in parsecs) from cluster center. Error-bars are computed as the Poissonian fluctuations of source counts in each bin.

tics on the number of stars in each annulus. We then assign a positive cluster identification if the radial profile exhibits at least two annuli with values above the background. In order to refine the location of the density peak, we repeat the radial density profile analysis starting from several locations within 10'' of the peak derived from the density maps; the location which maximizes the overall statistical significance of the annuli is then assigned as the cluster center. Figure 1 shows the typical footprint of a cluster, where the stellar density is plotted as a function of distance r from the start location; the density has a maximum at $r=0$ and decreases until it reaches a constant value which is the average background/foreground stellar density.

There are two exceptions in this analysis. The first is for source Mol160. The K_s -band image shows clear stellar density enhancement in a semi-circular annulus which surrounds the northern side of the dense millimeter core (see Figure A.26 in the Appendix), which appears devoid of stars. This stellar density enhancement is coincident with the emission patterns visible in the mid-IR (Molinari et al. 2008b) so it is clearly a stellar population associated with the star forming region. Since the millimeter peak is at the center of symmetry of the semi-circular stellar distribution, we will consider this as the center of the cluster. This is just for completeness of reporting, since we cannot say if the low density of stars at the millimeter peak is an effect of extreme visual extinction or reflects an intrinsic paucity of NIR-visible forming stars, as the proposed extreme youth of the massive YSO accreting in its depth would seem to suggest (Molinari et al. 2008b).

The second exception is for source Mol8. The stellar density analysis shows two peaks which are coincident with two distinct dust cores (see Fig. A.2); we then assume the presence of two distinct clusters, rather than a sub-clustering feature within the same cluster. The radial density profile analysis cannot be used here, so we fit elliptical gaussians to the peaks in the density maps, allowing for an underlying constant level representing the background stellar density. The resulting cluster richness is obtained integrating the fitted gaussian, and the cluster radius is taken equal to the fitted FWHM (the fitted gaussians are nearly circular).

Always following Testi et al. (1998), we determine the richness indicator of the cluster I_c by integrating the background-subtracted density profile; the cluster radius is taken as the radial distance from start location where the density profile reaches a constant value. This richness indicator is a very convenient figure to use in these cases where no detailed information is available for each single star in the region and the membership of the cluster cannot then be established for each single star. These values are reported in Col. 3 of Table 2 for all fields where a cluster has been clearly revealed. Col. 1 gives the target name (cf. Table 1); its kinematic distance is listed in Col. 2. The parameter N_{obs} (Col. 4) is the number of cluster members derived (see Sect. 4.1 below) from the integration of the background-subtracted K_s Luminosity Function (hereafter KLF, see Sect. 4.1). Also reported in Col. 8 is the mass of the hosting molecular clump; this is derived from the cold dust emission as reported in Molinari et al. 2008a, 2000), integrated over the entire spatial extent of the cluster; conversion into masses is done under optically thin assumption assuming $T=30$ K, $\beta = 1.5$ (Molinari et al. 2008a) and a mass opacity $\kappa_{230GHz} = 0.005 \text{ cm}^2 \text{ g}^{-1}$ which incorporates a gas/dust weight ratio of 100 (Preibisch et al. 1993). The IRAS source bolometric luminosity, Col. 9, is taken from Molinari et al. (1996, 2000, 2002, 2008a); in Col. 10 we list the A_V on the peak cluster position as estimated from submm observations (Molinari et al. 2008a, 2000). In Col. 11 and Col. 12 the coordinates of centers of the identified clusters are reported. Columns 6 and 7 contain parameters that will be described later in the text (see Sect. 3.2).

Following the described procedure, a cluster is detected within 1' of the IRAS position for 22 out of the 26 observed fields (85% detection rate). In two cases (Mol38 and Mol59) the stellar density map does not show a clear peak above the fluctuations of the field stellar density. For Mol 98 the radial density profile only shows one annulus above the background, so they fail the criterion that the stellar density enhancement should be significantly resolved above the background in two annuli. In one case (Mol30) several stellar density peaks are found in proximity of the IRAS source, but the lack of information of submillimeter/millimeter continuum prevents any firm conclusion.


Figure 2 shows the run of I_c as a function of peak A_V and suggests that with larger dust extinction it may be more difficult, or less likely, to detect a cluster at $2.2 \mu\text{m}$.

Our detection rate is quite high and tells us that young stellar clusters in sites of intermediate and massive star formation are essentially ubiquitous. While this evidence was established for relatively old Pre-MS systems like Herbig Ae/Be stars (Testi et al. 1999), we hereby verify that this is the case also in much younger systems where the most massive stars may even be in a pre-Hot Core stage (Molinari et al. 2008a).

Our detection rate is higher compared to other similar searches of stellar clusters toward high-mass YSOs. For example Kumar et al. (2006) use the 2MASS archive and report a rate of 25% (rising to 60% neglecting the inner Galaxy regions) toward a larger sample which also includes the sources of this work; in particular we detect all clusters also detected by Kumar et al. and in addition we reveal clusters toward 13 objects for which Kumar et al. report no detection. The reason for this discrepancy may be due to the fact that we obtained dedicated observations while Kumar et al. used data from the 2MASS archive; the diffraction-limited spatial resolution of our data is between a factor of 4 and a factor of 10 better with respect to 2MASS, and this certainly facilitates cluster detection especially in particularly crowded areas like the inner Galactic plane. To test this hypothesis we de-

Table 2. Results for Cluster Detection

Sou. Mol	d^a kpc	I_c	N_{obs}	R_{clu} pc	Pre-MS		M_{gas} $M_\odot$	L_{bol} $10^3 L_\odot$	A_V <i>peak</i> mag	Cluster Center	
					% CC	% CM				α (J2000)	δ (J2000)
3	2.17	78	78	1.7	34	99	910	12.4	18	00:44:57.4	+55:47:20.0
8A	11.5	25	30	1.3	37	27	1650	57.0	18	05:17:13.8	+39:22:29.7
8B	11.5	27	24	1.3	9	4	1780	5.5	8	05:17:12.0	+39:21:51.8
9	6.2	7	7	0.6	16	— ^g	—	24	—	05:20:16.9	+36:37:22.0
11	2.1	51	48	0.5	12	41	360	4.6	35	05:37:47.7	+31:59:24.0
12	1.6	12	13	0.3	0	30	72	1.6	46	05:40:24.4	+23:51:54.8
15	1.5	64	61	0.3	34	— ^g	—	5.8	—	06:08:41.0	+21:31:00.0
28	4.5	75	75	1.0	58	95	220	9.1	4	07:00:51.5	−08:56:18.2
30	0.3	200 ^e	no cluster detected ^b		—	—	—	0.14	—	—	—
38	0.5	−5	no cluster detected ^c		— ^h	—	—	0.19	40	—	—
45	4.3	28	27	0.5	68	19	1340	21.2	77	18:17:24.1	−17:22:12.3
50	4.9	46	43	0.6	36	41	80	17.3	25	18:19:07.6	−16:11:21.0
59	5.7	0	no cluster detected ^c		— ^h	—	—	11	29	—	—
75	3.9	8	7	0.7	37	24	1310	13.3	38	18:53:38.1	+01:50:26.5
82	6.8	8	10	0.3	0	52	590	15.4	52	18:59:03.2	+03:53:16.7
84	2.2	21	21	0.3	0	54	28	4.3	15	18:59:14.3	+07:04:52.3
98	4.5	6	no cluster detected ^f		— ^h	—	—	9.2	68	—	—
99	6.1	45	38	0.9	49	— ^g	—	37.3	—	19:11:51.4	+09:49:35.4
103	4.1	105	107	0.7	46	87	510	28.2	42	19:23:36.2	+17:28:58.1
109	4.3	19	17	0.5	^d	^d	1030	26.7	90	19:39:33.0	+24:00:21.3
110	4.3	23	20	0.5	^d	^d	400	14.8	55	19:40:58.5	+24:04:36.3
136	3.6	21	19	0.6	18	52	230	4	21	21:32:31.4	+51:02:23.1
139	7.3	25	24	1.2	10	31	1870	1.35	20	21:53:39.2	+56:27:50.7
143	5.0	25	22	0.8	10	76	630	7.8	26	22:19:09.0	+56:04:58.7
148	5.1	43	41	0.9	43	64	22	7.8	13	22:32:23.4	+58:19:01.3
151	5.4	15	14	0.9	12	30	2020	25	40	22:52:38.3	+60:00:44.6
160	5.0	36	34	1.3	30	76	1830	16	32	23:40:53.1	+61:10:21.0

^a Kinetic distance using the rotation curve from Brand & Blitz 1993.^b Stellar density analysis inconclusive due to extreme crowdedness of this field.^c Stellar density reveals no peaks close to the IRAS position or the submm peak.^d Only observed in K_s .^e Detection refused due to extreme field complexity (see text).^f Detection refused because only 1 annulus in the radial density profile is above background (see text).^g No extinction estimate is available due to lack of submm information to evaluate de-reddening correction.^h Extinction estimate is available from single-pointing submillimeter data (Molinari et al. 2000) but not from maps, so that a reliable clump mass estimate is not possible.**Fig. 2.** Cluster richness indicator I_c as a function of A_V on cluster center; for a few detected cluster we do not have an estimate of A_V .

graded the NTT K_s image of Mol103, also considered in Kumar et al., to the 2MASS resolution; extraction and photometry were performed as outlined above but the search for a cluster based on the stellar radial density profiles revealed no cluster. Besides, the estimated number of members (corrected for the contribution of fore/background stars) for 7 out of the 10 clusters detected both by us and by Kumar et al. is at least a factor of two less in the latter study.

Kumar & Grave (2008) conducted a similar study on a large sample of high-mass YSOs, that include a certain number of our sources, using this time data from the GLIMPSE survey (Benjamin et al. 2003). In this work they detect no significant cluster around any targets in a sample of 509 objects. As the authors say in the paper, however, GLIMPSE data are sensitive to 2–4 $M_\odot$ pre-main sequence stars at the distance of 3 kpc. Based on color-magnitude analysis (see later below) our mass sensitivity is of the order of 1 $M_\odot$ at a distance of 3.6 kpc and $\sim 0.6 M_\odot$ at a distance of 2.1 Kpc. Probing longer wavelengths, GLIMPSE is likely to be more sensitive to younger sources compared to the classical J, H, K range which also samples relatively older pre-MS objects. The combination of sampling higher-mass (and hence more rare stars due to the shape of the IMF) and relatively younger stars (which, as indeed our analysis finds, may not be

the majority in a young cluster) may plausibly be the reason of the negative cluster detection results in Kumar & Grave.




Fig. 3. Distribution of the cluster's radii in parsecs (full line), and of the cluster richness indicator I_c in number of stars (dashed lines); the median values for the two distributions are 0.6 pc and 37 stars, respectively

The distribution of the radii of the detected clusters is shown with the full line in Fig. 3; the median value is 0.7 pc. The dashed histogram (which refers to the upper X-axis) shows the distribution of the cluster richness indicator I_c , with a median number of stars of 27. We note that the value of I_c for many of our clusters is less than the limit of 35 suggested by Lada & Lada (2003) to be a *bona fide* cluster. This definition stems from the argument that a less rich agglomerate may not survive the formation process as an entity. Our interest, however, is to investigate the spatial properties of the young stellar population in a star forming region at the time of active formation, without worrying about its possible persistence as a cluster at the end of the formation phase. However, we prefer not to introduce a new term to identify the structures we see and still use the term cluster, although in a "weaker" sense compared to Lada & Lada.

3.2. Properties of Identified Clusters

We will first derive qualitative indications concerning the nature of the identified clusters using simple diagnostic tools like color-color and color-magnitude diagrams. These diagrams have been drawn for all detected clusters and are available in electronic form; we here illustrate the particular case for Mol28.

3.2.1. Color-Color Analysis

Fig. 4 shows the $[J-H]$ vs $[H-K_s]$ diagram for all sources detected within a distance equal to R_{cl} centered on the stellar density peak. The full circles represent all sources detected in all three bands, the arrows represent sources with lower limits (in magnitude) in the J band. The plot shows more stars than the I_c value reported in Table 2 because we also include the fore/background stars that cannot be individually identified against the true cluster members. A significant fraction of the sources have colors compatible with main-sequence stars with a variable amount of extinction reddening (computed adopting the Rieke & Lebofsky (1985) extinction curve), but many sources show colors typi-

cal of young pre-MS objects with an intrinsic IR excess arising from warm circumstellar dust distributed in disks (Lada & Adams 1992). The set of dotted curves represents the locus of two-component black-bodies with temperatures as indicated at the start and end of each dotted line; along each curve the relative contribution of the two blackbodies is varied. These curves mimic the effect of a temperature stratification in the dusty circumstellar envelopes, and the presence of sources in the area covered by these curves is an indication of the presence of warm circumstellar dust.




Fig. 4. $[J-H]$ vs $[H-K_s]$ diagram for Mol28. $[J-H]$ is obviously a lower limit for sources not detected in J. The continuous curve at the bottom-left represents the Main Sequence, while the dashed grey lines represent the effect of reddening (Rieke & Lebofsky 1985) for variable amounts of extinction as indicated along the lines. The dashed-dotted black line is the Black-Body curve, and the dotted curves are two-component Black-Body with varying relative contribution (respectively, from the inner to the outer curve, 3000-1500K, 3000-1000K, 3000-900K and 3000-500K).

A straightforward indication of the youth of the cluster may be offered by the fraction of sources which are not compatible with reddened MS stars, i.e. those with IR excess. The number of stars with IR excess is normalized to the total number of stars revealed in the cluster area corrected for the expected number of fore/background stars estimated from the areas surrounding the cluster (but still in the same imaged field). To be conservative we extend the region of the MS by 0.2 magnitudes to the right corresponding to about 2σ uncertainty on measured magnitudes. This ratio is reported as a percentage value in Col. 6 of Table 2).

3.2.2. Color-Magnitude Analysis

Additional evolutionary indications for the detected clusters may be derived from the K_s - $[H-K_s]$ diagram, reported for Mol28 in Fig. 5. Compared to the main sequence (the leftmost almost vertical curve in the figure) a significant fraction of the sources are on its right, where the evolutionary tracks for Pre-MS sources (Palla & Stahler 1999) can also be found, and could be therefore interpreted as very young pre-MS objects. The distribution of sources in the diagram spans a much larger region than the one covered by the Pre-MS isochrones, and this is due to the combined effect of extinction reddening and IR excess. Extinction effects can be appreciated looking at the dotted lines originat-

ing on the main sequence and extending toward bottom-right for increasing values of A_V ; on the other hand the presence of a warm dusty circumstellar envelope implies an increase in absolute emission and in SED steepness, therefore shifting a pure photosphere toward top-right of the diagram (as shown by the arrow labeled 'IREX' in Fig. 5). Similar to the color-color analysis, it is impossible to try and estimate the age of individual stellar sources based on their location on the pre-MS isochrones, because we do not know the amount of A_V by which we should deredden each object. We follow a conservative approach by dereddening each object using half of the extinction estimated for each location from millimeter maps; this corresponds to putting each object midway through the clump.




Fig. 5. K_s vs $[H-K_s]$ diagram for Mol28. The leftmost curve represents the Main Sequence, while the dashed lines represent the effect of reddening for variable amounts of extinction. Isochrones from Palla & Stahler (1999) are also indicated with full lines for different Pre-MS ages. The arrow labeled IREX indicates the direction of change due to IR-Excess (see Sect. 4.2). $[H-K]$ is obviously a lower limit for those sources not detected in H. Symbols in grey color indicate sources with IR excess as determined from the color-color diagrams (see fig. 4).

A further correction is to remove the IR excess for those sources in which this is apparent in the color-color diagram (see fig. 4), as estimated using the formulation suggested by Hillenbrand & Carpenter (2000), and that will also be used later in this work (see Sect. 4.2). The fraction of pre-MS stars over the total in each cluster area will still be contaminated by foreground/background stars; to estimate this contamination we choose an off-cluster area in the same imaged field and we simply compute the ratio of sources with pre-MS colors over the total (in these off-cluster regions there is no significant reddening to correct for). Col. 7 of Table 2 reports for each cluster the fraction of stars (detected in the cluster area in all three bands) situated more than 0.2 mag to the right of the MS after the various corrections have been applied.

4. Initial Mass Functions and Star Formation Histories

As it is apparent from the qualitative analysis presented in the previous paragraphs, the diagnostic power of our observations is limited because we do not know which objects in the cluster area

are real cluster members and we do not know the exact amount of dust extinction (originating within the hosting clump) and IR excess (originating in the immediate circumstellar environment) pertaining to each source. Lacking the detailed knowledge on individual stars in the clusters, fundamental quantities like the Initial Mass Function (IMF) and the Star Formation History (SFH) cannot be directly derived from, e.g., the K_s luminosity function (KLF). We are then forced to obtain these using statistical simulations of clusters based on different input parameters and performing a statistical comparison between synthetic and observed KLFs and HKCFs.

We will first derive the observed KLFs from the observations; we will then illustrate in detail the model used for the cluster simulations, exploring the sensitivity of the results to a wide range of input parameters; finally, modeled and observed KLFs will be compared to infer statistical indications for the IMF and SFH for our clusters.

4.1. Observed K_s Luminosity Functions

The KLF for each cluster is obtained simply counting all detected sources within the cluster area as identified from the cluster density profile (see §3.1). Similar to the other diagnostic tools (§3.2.1 and 3.2.2), the KLF will be contaminated by field stars that cannot be individually identified. To account for the field star contamination in a statistical way we subtract from the KLF built on the cluster area, the KLF built in a region outside the cluster area but still in the same imaged field, after normalising for the different areas. Regions where the field star KLF is built have a lower extinction respect to the cluster one, so the background contribution to the cluster KLF is likely to be overestimated. Field-subtracted KLFs for all clusters are presented in Appendix 6, and are also available online¹.

The integral of the KLF gives an independent estimate of the number of cluster members; these values are reported as N_{obs} in Table 2; their agreement with richness indicator I_c confirms the consistency of the analysis. All KLFs show a dominant peak which always lies close to the completeness limit, showing that our observations are not sensitive enough to trace the low-mass stellar component of our clusters. Many of the KLFs present a separate small peak at low magnitudes (one or two sources at most, on average). Can this be due to confusion arising from source crowding and insufficient spatial resolution? We studied for each cluster the distribution of distances of each star from its nearest neighbour and we find that there are essentially two types of distributions, reported in Fig. 6. In the first type (full line in figure) the distribution has a peak corresponding to an inter-star distance significantly higher than the value corresponding to half the PSF FWHM (the full vertical line); in this case the suggestion is that all cluster members have been resolved from their neighbour. In the second type (dashed line in the figure) the distribution has its peak very close to half the PSF's FWHM (the dashed vertical line), indicating that source blending should be certainly considered possible. We verified that all clusters exhibiting a distance distribution of the second type do show a second faint peak at high brightness in their KLFs, therefore confirming that this feature is an artifact of the relatively low spatial resolution which in some cases is not sufficient to resolve all cluster members.

¹ at <http://galatea.ifs-roma.inaf.it/faustini/KLF/>




Fig. 6. Distribution of identified sources as function of nearest-neighbour distance (D) for two of examined fields (Mol28 dashed line and Mol103 full line).

4.2. Synthetic KLF. Synthetic Cluster Generator: a Near-IR Cluster Simulator

As already mentioned, from our data alone we cannot derive masses and ages. We thus developed a model to create statistically significant cluster simulations obtained for different assumptions of IMF and SFH (source ages and their distribution), and compare the synthetic KLFs with the observed field-subtracted KLFs. This model we call Synthetic cluster Generator (SCG).

4.2.1. SCG: Model Description

A cluster is created by adding stars whose masses and ages are assigned via a Monte-Carlo extraction according to the chosen IMF and SFH; the pre-MS evolutionary tracks of Palla & Stahler (1999) are then used to convert them into J , H and K_s magnitudes. The 3D distribution of stars is obtained by randomly choosing for each star a set of x, y, z coordinates using the observed stellar density profile (see Sect. 3.1), approximated as a radially symmetric Gaussian, as weight-function; this is needed to assign, using submm continuum images, the proper column of cold dust to extinguish the near-IR radiation. Other analytical functions could have been used, e.g. a King profile, but the statistics of our clusters are not sufficiently high to try and explore the effect of different radial profile assumptions.

To convert the submm flux into dust column density we used the dust temperature and emissivity exponent β as determined in Molinari et al. (2000); mean values from the latter work were adopted for those fields not covered as part of that work.

To properly simulate the pre-MS stars, we also need to include the effect of an IR excess due to warm dust in the circumstellar envelopes and disks. We used the distribution (modeled as a Gaussian) of $[H-K_s]_{ex}$ color excesses as measured for a sample of Pre-MS stars in Taurus, as used by Hillenbrand & Carpenter (2000), as a weight-function to randomly assign a $[H-K_s]_{ex}$ to each simulated star in our model; the K_s vs $[H-K_s]_{ex}$ relationship adopted in the above mentioned work was then used to derive the H and K_s excess-corrected magnitudes. The K_s magnitude of the synthetic star was then compared with the limiting magnitude typical of the cluster being simulated to determine if the star could have been detected in our observations. This procedure is

repeated until the number of synthetic detectable stars equal the value of I_c as determined in our observations; at this point the cluster generation process is complete.

Since the simulation is based on Monte-Carlo extraction for stellar mass, age and position in the cluster, each independent run for a fixed set of input parameters can in principle result in very different outputs in terms of cluster luminosity, total stellar mass, maximum stellar mass and synthetic KLF. To provide statistical significance, the model is run 200 times for any given set of input parameters, and the median KLF is later adopted for the comparison with the observed one. Clearly, the predictive power of this simulation model resides in its capability to characterize the cluster properties for any given parameter set. In other words, the distribution of the resulting quantities should not be uniform but peaked around characteristic values. We will come back to this in section Sect. 4.2.3

4.2.2. SCG: Input Assumptions

We tested three different assumptions for star formation histories in our cluster simulations. The first choice is to assume that stars in the cluster formed in a single burst-like event (hereafter SB) some t_1 years ago. The explored range in the simulations is $10^3 \leq t_1 \leq 10^8$ yrs. The second choice is to have the formation of stars proceed at a constant rate (hereafter CR) from a time t_1 years ago to a time t_2 years ago. The ranges explored in the simulations are $10^4 \leq t_1 \leq 10^8$ yrs and $10^3 \leq t_2 \leq 10^7$ yrs, where always $t_1 > t_2$. The third possibility we explored is a variation of the previous one, where the star formation rate is not constant but varies with time in a Gaussian fashion (hereafter GR). Within the boundaries for start and end of the star formation process, t_1 and t_2 varied as above, we also varied both the time t_c for the peak of the Gaussian in the range $10^{3.7} \leq t_c \leq 10^{7.7}$; the $\text{Log}_{10}(\sigma)$ of the Gaussian-like SFH was allowed to assume the two values 0.1 and 0.5.

As for the IMFs, we allowed three different choices from Kroupa et al. (1993), Scalo (1998) and Salpeter (1955), with the latter modified introducing a different slope for $M < 1 M_\odot$ that coincides with that of the Scalo (1998) IMF; the three IMFs will be coded as IMF1, IMF2, and IMF3 respectively. The IMF from Kroupa et al. favors the low-mass end of the distribution, while the classical Salpeter IMF is flatter at low mass but heavier at intermediate and high masses (above $1 M_\odot$). The IMF from Scalo is kind of intermediate between the two, resembling Salpeter's one below $1 M_\odot$ and above $10 M_\odot$, and Kroupa's for $1 M_\odot < M < 10 M_\odot$.

4.2.3. SCG: Predictive Power

In order to verify our model's predictive power, we ran a set of 200 simulations for a cluster with a Salpeter IMF and a constant star formation rate with $t_1 = 10^6$ yrs and $t_2 = 10^4$ yrs. Figure 7 shows the distribution of the predicted number of stars and the total luminosity for the 200 simulations. The number of cluster members shows very little variations, as expected since the number of detectable stars is the parameter we use to stop the simulation; on the other hand the distribution of the total luminosity is not particularly peaked, as the central 3 bins containing about 60% of the simulations span almost two decades in luminosity.

On the other hand the distributions for the total cluster stellar mass, and for the mass of the most massive member (see Fig. 8) are rather peaked and highlight a relatively higher predictive power of the model for these two quantities. It is to be




Fig. 7. Distribution of the predicted number of cluster members (full line) and total luminosity (dashed line) for 200 SCG runs for Mol160 with a Salpeter IMF and a constant star formation rate with $t_1=10^6$ yrs and $t_2=10^4$ yrs.

noted that the distributions are rather skewed, suggesting that neither the mean nor the median are particularly suited to characterize the peak of the distribution. Indeed we have computed the values that these quantities assume at the peak of their distributions, to have a more representative value for the masses and use them in the following arguments.




Fig. 8. Distribution of the predicted total stellar mass (full line) and mass for the most massive star (dashed line) in a cluster for 200 SCG runs for Mol160 (same inputs as in Fig. 7).

Concerning the reproducibility of the KLF, for each of the 200 runs the resulting KLF was fitted with a Gaussian function and the center, peak and σ were determined. Fig. 9 reports the distribution of these three parameters for the 200 runs and shows that all of them are remarkably peaked and symmetric. The formal r.m.s. spread for the three quantities, estimated via a Gaussian fit to the distributions in the figure, is ≈ 0.3 mag for the KLF center, $\approx 12\%$ for the KLF peak (about 1.2 sources out of a mean KLF peak of 10), and ≈ 0.25 mag for the KLF FWHM.




Fig. 9. Distribution of the predicted center magnitude (full line - bottom X-axis scale), width (dotted line - bottom X-axis scale) and peak value (dashed line - top X-axis scale) of the predicted Gaussian-fitted KLFs for 200 SCG runs for Mol160 (same inputs as in Fig. 7).

We have made a similar analysis for HKCF (H-K_s color function; see Sect. 3.2.2). Figure 10 shows the distribution of Gaussian function centers, peaks and σ 's for HKCFs obtained for the same 200 runs used previously for the KLFs. Gaussian fits to the three distributions in the figure give an r.m.s. that is ≈ 0.15 mag for the HKCF center and ≈ 0.14 mag for the HKCF FWHM, while the 'peak' distribution is flatter and has an r.m.s. value of $\approx 21\%$ for the HKCF peak (about 3.2 sources out of a mean HKCF peak of 15). It is worthwhile to stress that since the position which is assigned to each simulated star in the cluster is different in each of the 200 runs of the model (for any given set of input parameters), the scatter in the properties of the synthetic KLFs and HKCFs also statistically tends to account for the effects of extinction variations in the cluster's hosting clump, which may in principle be relevant in such heavily embedded systems (see Tab. 2).




Fig. 10. Distribution of the predicted center color (full line - bottom X-axis scale), width (dotted line - bottom X-axis scale) and peak value (dashed line - top X-axis scale) of the predicted Gaussian-fitted HKCFs for 200 SCG runs for Mol160.

We conclude that the model results, for a given set of input parameters, have a good reproducibility, except concerning the total luminosity. The model therefore has a quite fine predictive power concerning the median properties of a synthetic cluster. Indeed, the spread of KLF center magnitudes is less than the bin amplitude used in building the KLFs for the simulations (and that will be used for the rest of the work); the median synthetic KLF therefore provides a good representation of the cluster luminosity distribution.

In conclusion, 200 simulation runs for each combination of input parameters (IMF and SFH) seem enough to achieve a sufficient statistical significance for the properties of the synthetic observables (KLFs and HKCFs). Although the distributions for the KLFs' (HKCFs') parameters seem rather symmetrical, we will adopt the median KLF (HKCF) of the 200 runs as a more reliable characterization for that particular parameters' set. Using the mean KLF (HKCF) for the comparison does not significantly alter the results.

4.2.4. Exploring the SCG parameter space: Cluster Parameters

After verifying the robustness of model results for independent runs for the same input parameters, we now want to verify the sensitivity of the model results against changes of these parameters. We will first concentrate on simulated cluster physical parameters (number of cluster members, total luminosity, stellar mass distribution), and in the next paragraph we will examine how the KLFs and the appearance of the color-magnitude diagrams, which are the main observables we will use in our analysis, behave in this respect.

Number of stars N_{stars} - As a general rule the older the cluster is allowed to be, irrespective of the detailed SFH adopted, the higher is the number of produced stars. This is easily understood since the SCG cluster building is stopped when the number of the K_s -detectable stars equals the number of observed objects; if a cluster is old the stars will be intrinsically fainter due to the shape of Pre-MS tracks, and it will statistically be less likely to extract stars bright enough to be detectable. N_{stars} does not significantly depend on the IMF choice as long as $t_1 \leq 10^6$ yrs, while for older systems IMF1 (Kroupa et al. 1993) will produce nearly twice as many stars as IMF3 (Salpeter 1955) with IMF2 (Scalo 1998) in between.

Stellar Masses - Likewise, the total stellar mass and the mass of the most massive star will be higher the older the cluster is allowed to be. If an IMF1 cluster is a very old SB or a CR with $t_1=10^8$ yrs and $t_2=10^7$ yrs for example, $M_{\star Tot}$ and $M_{\star Max}$ will be respectively a factor of 5 and 2 higher with respect to clusters which are younger and/or are allowed to form stars until recent times (i.e. allowing a CR with $t_2 = 10^4$ yrs). The explanation follows directly from the argument made for the N_{stars} behavior above; matching the number of K_s -detected stars in a relatively old cluster with intrinsically fainter stars will require that stars will have to be on average more massive objects, and this will clearly result also in a higher total stellar mass.

Going from IMF1 to IMF3 both $M_{\star Tot}$ and $M_{\star Max}$ will significantly increase, as expected. The trend of $M_{\star Tot}$ with cluster age is less pronounced because with IMF2 and IMF3 it is statistically more likely to produce relatively more massive (and hence more easily detectable in K_s) stars requiring a lower num-

ber of star extractions and hence a lower relative total mass at the end of the simulation. The age-trend of $M_{\star Max}$ is instead the same (only shifted toward higher masses) because the probability of extracting a massive star is the same for all ages and is only function of the chosen IMF.

Total Stellar Luminosities and Massive Object Luminosities - The total stellar luminosity, like the luminosity of the most massive star ($L_{\star Max}$), exhibits the same behaviour as $M_{\star Max}$. This is easily understood given the steep power-law dependence of the stellar luminosity on mass, and confirms that the total luminosity (L_{Tot}) will be largely dominated by the most massive stellar object in the cluster: $L_{Tot} \propto L_{\star Max}$. Of great interest is the ratio between $L_{\star Max}$ and L_{Tot} ; for the large majority of clusters its value varies between 0.6 and 0.8. This is further confirmation that global properties of our clusters are dominated by the most massive source. This ratio does not present any particular dependence on the value of $M_{\star Max}$, or of N_{stars} ; only for the most populated clusters (clusters with of the order of a hundred members, such as Mol103, where the contribution of a great number of low-mass sources becomes important, do we find a lower value for this ratio.

4.2.5. Exploring the SCG parameter space: KLF variations

We will now briefly analyze the diagnostic power of the KLF and the HKCF against changes in IMF and SFH choices. Figure 11 shows the KLFs predicted for source Mol3 adopting the same SFH parameters (as indicated in the figure) and using the three different IMF choices.




Fig. 11. KLF (using the absolute K_s magnitudes) for Mol3 predicted by SCG for a CR cluster with $t_1=10^7$ yrs and $t_2=10^4$ yrs, for three different choices of the IMF (line styles as indicated in the figure). The dash-dotted line represents the completeness limit for this source given the K_s limiting magnitude.

The shape of the resulting KLF changes throughout the M_K range; going from Kroupa et al.'s IMF1 to Salpeter's IMF3 the distribution gets more skewed toward lower magnitudes; this was expected since IMF1 produces more lower mass stars than IMF3. One can certainly argue that the change is not dramatic, but on the other hand the modification does not affect one or two bins but the entire KLF consistently. The change is more appar-

ent in the region between the peak and the completeness limit than at the bright end of the KLF, and for this reason the ability of the model to discriminate among different IMFs will be better for those sources, as Mol3 in the figure, where the KLF's peak is clearly detected above the completeness limit.




Fig. 12. KLF (using the absolute K_s magnitudes) for Mol3 predicted by SCG for an IMF3 cluster with different ages as indicated in the plot. Older clusters produce a KLF peaked toward lower magnitudes. The dash-dotted line represents the completeness limit for this source given the K_s limiting magnitude.

The difference in predicted KLFs is much more dramatic if different age ranges are assumed, keeping fixed the shape of the SFH and the IMF, as it is apparent in Fig. 12. The peak of the KLF shifts considerably toward higher magnitudes as the median stellar ages (t_c) increase. A similar trend is observed comparing SB models with different ages, although SB models always produce KLFs which are considerably narrower than CR or GR models. Larger cluster ages would shift the peak of the KLF beyond the completeness limit; in other words, our analysis is not sensitive to ages for the majority of stars in excess of $\sim 5 \cdot 10^6 \div 10^7$ yrs; besides, such old clusters would be hard to justify given the fact that they are still heavily embedded in dense clumps.

Finally, we briefly show how the KLFs change for different choices of the SFHs. Figure 13 shows the KLFs obtained for a SB with $t_1=10^6$ yrs, compared with a CR with $t_1=10^7$ yrs and $t_2=10^4$ yrs and a GR with the same start and end star formation period, and with a peak times for star formation rate of $t_c=10^6$ yrs. The KLFs are clearly different, with a peak magnitude which is quite sensitive to the formation rate typology and the peak time for star production.

As concerns the $[H-K_s]$ color functions, they are found insensitive to the choice of IMF. Similarly to the case of the KLFs, the main differences between synthetic HKCFs are more evident for different age ranges especially in the number of detectable stars.

4.3. Comparing Observed and Synthetic KLFs and HKCFs

The detailed comparison of the model KLFs and HKCFs functions to those observed was carried out only for those sources where the number of detected stars was sufficient ($I_c \geq 15$) to allow a statistically significant comparison (see Col. 3 of Table 2),




Fig. 13. KLF (using the absolute K_s magnitudes) for Mol3 predicted by SCG for an IMF3 SB cluster with $t_1=10^6$ yrs (full line), CR with $t_1=10^7$ yrs and $t_2=10^4$ yrs (dotted line), and GR with same t_1 and t_2 and $t_c=10^6$ yrs (dashed line). The dash-dotted line represents the completeness limit for this source given the K_s limiting magnitude.

and where submm information was available to allow meaningful estimates of extinction (this excludes Mol15 and Mol99). The number of clusters fulfilling these criteria were 16 out of 23 detected clusters. The comparison of the observed KLFs and HKCFs (KLF_{Obs} , $HKCF_{Obs}$), with the synthetic ones produced by SCG (KLF_{Syn} , $HKCF_{Syn}$), for the full set of input parameters (IMF, SFH and age parameters) was carried out automatically; to ease the process, the observed and synthetic functions were computed on the same M_K and H-K grid.

The comparison procedure between synthetic and observed KLFs is described below, but it is the same for HKCFs. The KLFs are first compared bin by bin (the comparison being limited to those bins brighter than the completeness limit) identifying with i each bin of the observed KLF, starting from $i=1$ for the lower- M_K non-zero bin to N for the bin where the completeness limit for that source is reached (the number N will clearly be different for each cluster). In the case of HKCFs, we exclude objects with H and K magnitudes greater than the observed limiting magnitudes for these bands. A matching flag m_i is set to 1 for those bins where the number of sources coincide within the 1σ Poissonian error bar of the observed KLF, i.e.:

$$|N_{\star Syn_i} - N_{\star Obs_i}| \leq \sqrt{N_{\star Obs_i}} \quad (1)$$

The total number of bins where a match is found is divided by the total number of bins useful for the comparison to get a KLF compatibility figure (in %) as

$$C = 100 \times \frac{\sum_{i=1}^N m_i}{N} \quad (2)$$

The higher is C , the better is the overall match between KLF_{Obs} and KLF_{Syn} .

However, the same value of C may result from bins concentrated in the low- M_K end of the KLF, where there are few sources, or in the region around the peak and in the proximity of the completeness limit, where instead there are more sources and

Table 3. Results for SCG runs on detected clusters

Source Mol	IMF	$t_{10\%}$	t_{med} yrs	$t_{90\%}$	$N_{\star}$	$M_{\star Max}$ $M_{\odot}$	$M_{\star Tot}$ $M_{\odot}$
3	1	$10^{4.98}$	$10^{5.40}$	$10^{5.81}$	91	3.7	61
8A	2-3	$10^{4.71}$	$10^{5.38}$	$10^{5.83}$	31	3.8	30
11	1-2-3	$10^{6.06}$	$10^{6.41}$	$10^{6.82}$	48	3.8	47
28	3	$10^{6.03}$	$10^{6.12}$	$10^{6.24}$	77	9.9	105
50	1	$10^{6.16}$	$10^{6.48}$	$10^{6.66}$	53	3.5	36
103	1-2	$10^{6.57}$	$10^{6.70}$	$10^{6.83}$	115	4.0	80
139	3	$10^{5.34}$	$10^{5.96}$	$10^{6.46}$	25	2.9	16
143	2	$10^{6.57}$	$10^{6.70}$	$10^{6.82}$	27	3.1	21
160	1-2-3	$10^{6.57}$	$10^{6.7}$	$10^{6.83}$	89	4.3	63

hence higher statistical significance. A further figure of merit is then introduced,

$$W = \sum_{i=1}^N \frac{m_i \cdot N_{\star Obs_i}}{N_{\star Obs_{tot}}}, \quad (3)$$

where $N_{\star Obs_{tot}}$ is the total number of sources present in all the bins used for comparison. This parameter weights each matching bin by its relative richness, favoring the bins closer to the completeness limit and the KLF peak over those in the bright tail of the distribution. This choice is due to the fact that the KLF (HKCF) peak region is the most sensitive to changes in the SCG input parameters.

In this automatic procedure we select those models for which the parameters C and W (at the same time for KLFs and HKCFs) are maximum. For Mol8B, Mol45 and Mol84 the observed KLF has a very irregular and multiple-peaked shape which cannot be matched by any model, and are therefore discarded from further considerations. We are then left with 14 clusters for which a series of models can be found with at least 75% of the bins matching the observations. We find that the best values of C and W are never found for one single set of parameters, but rather we identify ranges of parameter values which produce the best match; in other words there is a level of degeneracy that the models cannot remove, and this varies from source to source. In 4 clusters (Mol109, 110, 136 and 151) this degeneracy is essentially complete and the model is not able to make any prediction; in one case (Mol148) the comparison selects models with very old stellar ages but with total stellar luminosities by far in excess of the measured bolometric luminosity obtained integrating the integrated observed for this region from the mid-IR to the millimeter (see Tab. §2). In the 9 remaining cases some degeneracy persists especially in the IMF, confirming (§4.2.5) that our models are weakly selective on the IMFs, but there are clear indications concerning the SFH and ages.

Table 3 reports a summary of the results. The IMF of matching classes of synthetic cluster models is shown in Col. 2. Cols. 3, 4 and 5 contain the times for the formation of 10%, 50% and 90% of the total number of cluster members; these values are obtained as the median of the values that these times have in all models that match the observations. Col. 6 is the number of cluster members $N_{\star}$; Col. 7 shows the mass of the most massive object $M_{\star Max}$; Cols. 8 reports the total stellar mass of the cluster $M_{\star Tot}$. We stress again that the analysis selects classes of models rather than single models; the values reported in Table 3 are the median of the values that the parameters assume within the class of matching models. The table shows that for some

fields multiple IMFs are compatible with the data and, in general, SFHs with constant (CR) or Gaussian (GR) star formation rate provide acceptable solutions for certain age ranges (as reported in the Table). Simulations with SFHs with a single burst are, in general, not accepted. Our modeling is insensitive to bulk stellar ages in a cluster in excess of $5 \cdot 10^6 \div 10^7$ years (§4.2.5).

5. Discussion

5.1. Cluster Ages and Star Formation Histories

Perhaps the most important result of this work is that in all clusters where the comparison of observed KLFs with the ones predicted by the SCG model is possible (see previous paragraph), the observations are consistent with a star formation which goes on over time intervals that in most cases have a spread between about few 10^5 and few 10^6 yrs, and with a median cluster age of a few 10^6 yrs. In most cases we cannot discriminate clearly between a constant or variable SFR but we are confident that we can exclude that on average the stars in our clusters are co-eval and originating from a single burst of formation. Detailed studies toward the Orion Nebula Cluster show that stars have been forming for at least $10 t_{dyn}$, or $20 t_{ff}$ (Palla & Stahler 1999; Hoogerwerf et al. 2001, Hillenbrand 1997), and our results would seem to generalize this on a larger sample of intermediate and high-mass star forming regions.

In principle it can be argued that our analysis is incomplete since we did not take in consideration longer wavelength data which could pinpoint heavily extincted objects barely visible, or not visible at all, in the near infrared. This however, does not appreciably modify our conclusions about the age spread within the clusters. Indeed, Vig et al. (2007) applied a different analysis to the specific region Mol075, a field not included in our final analysis (Table 3) because the background-subtracted KLF is populated by too few objects for a statistically significant model comparison. Vig et al. also considered *Spitzer* IRAC and MIPS data, looking for the brighter and redder objects in the area covered by submillimeter emission. In this way they could identify the younger and more massive objects in the field with an estimated age of the order of 10^6 years or less. This approach, however, is not sensitive to low mass and relatively older pre-MS objects, for which our method is ideally designed. While for this particular field, for the reasons explained above, we cannot perform a direct comparison to our approach, it is clear that the inclusion of longer wavelength data in the analysis might have identified a different, younger, population of objects, rather amplifying the observed age spread deduced for the clusters.

Models of cluster formation via competitive accretion seemingly succeed in delivering an IMF close to the observed ones thanks to the spread of the accretion rates consequent to the competitive accretion mechanism, but the prediction that all stars are formed in about 5×10^5 yrs (Bonnell et al. 2004) for typical conditions in young clusters, corresponding to a dynamical time or so, seems to be in disagreement with our results. We instead favour scenarios (Tan & McKee 2002) where stars keep forming over several free-fall times thus providing the required age spread. The finding that the most massive object in the fields considered in this work are still being formed or have just finished a phase of intense accretion (Molinari et al. 2008a) is a further indication that star formation **seems** to be a long-duration process in the life of a molecular clump.

How do our clusters compare to relatively more evolved system, like those observed for a sample of Herbig Ae/Be by Testi et al. (1997, 1998) ? Figure 14 shows the relationship between the




Fig. 14. Number of cluster members as a function of Mass of highest mass star. Asterisks are for Testi et al.'s (1997, 1998) Herbig Ae/Be sample. Other point are our source sample, where the full diamonds are the N_* obtained from our clusters simulation analysis; the full lines passing through the full diamonds represent the total spread of the parameters from all SCG models that match the observed KLF and HKCF. Full triangles are the same clusters where this time we use the observationally derived I_c instead of the model-provided N_* . Empty triangles are those clusters which were not analyzed with SCG, or which exhibit complete model degeneracy; in these cases I_c was used for the cluster membership (from Table 2), while the maximum stellar mass has been derived assuming that half of the bolometric luminosity is generated by a single ZAMS star (the horizontal lines through the empty triangles represent the mass spread assuming that a fraction between 30% and 100% of L_{bol} come from a single star)

mass of the most massive source and the total number of cluster stars as provided by the SCG simulations for our modeled clusters. The filled symbols represent the clusters which we could model (Tab. 3); the empty symbols represent the clusters which instead could not be modeled for a variety of reasons (see §4.3), while Testi et al.'s clusters are reported as asterisks (see figure caption for detailed explanation of the symbols). The figure suggests that the clusters presented in this study are richer than those surrounding Herbig Ae/Be stars for any given value of the most massive star in each cluster. The trend persists if we use similar indicators (e.g. I_c , the full triangles in the figure). Furthermore, we note that while the limiting magnitudes of our observations and those of Testi et al. (1998) are similar, higher A_V values toward our sources and the typically greater distance from the Sun would justify the non detection of the fainter cluster members predicted by the SCG models. It is thus likely that the values of I_c derived from our observations tend to underestimate the cluster memberships.

This evidence is clear for values of the most massive star in the cluster below $\sim 10M_\odot$, where the 9 clusters for which we could compare observations with SCG predictions lie (the diamonds). In the clusters for which this could not be done (the empty triangles), an estimate of the mass for the highest-mass star was made by assuming that a fraction between 30 and 100% of the bolometric luminosity is due to a single ZAMS star. In this case the trend for richer clusters compared to Herbig stars (i.e., the asterisks) would tend to become marginal. These estimates, however, place the latter clusters systematically to the right in the plot, compared to the 9 modeled clusters; indeed, if we were to estimate in the same way a maximum stellar mass also for the 9 modeled clusters, we would obtain values in excess (between a factor two and three) to those provided by the detailed SCG modeling. In other words, the evidence that our clusters are richer than those around Herbig stars is marginal at worst (i.e., using the most conservative approach of estimating the mass of the highest-mass star).




Fig. 15. Distribution of radii for our clusters (full line) and those associated with Herbig Ae/Be stars (from Testi et al. (2001), dashed line).

The plausibility of this interpretation is strengthened by the recent results of Baumgardt & Kroupa (2007) who carried out extensive numerical simulations of the evolution of stellar clusters as a function of, among other parameters, the cluster gas content. They show that for a wide range of initial conditions and star formation efficiencies, the dispersal of the gas with age causes the cluster to expand overall and disperse a fraction of the stars originally belonging to the cluster. Besides, as the cluster expands its decreasing stellar density would make the low-density outer regions of the cluster ever more difficult to detect against the field stars (especially in the Galactic plane, where all these objects lie), mimicking a smaller cluster from an observational viewpoint. Fig. 15 shows the distribution of radii for our clusters (full line) and for those surrounding Ae/Be stars; radii have been derived with the same analysis in the two samples and the histogram clearly shows that our clusters are indeed larger in size, confirming the prediction of Baumgardt & Kroupa (2007).

This age effect on clusters size is also revealed within our clusters sample. Fig. 16 presents the relationship between the clusters radii as derived from the observations and their ages as derived from the modeling. The reported correlation has a Spearman rank correlation coefficient of ~ -0.6 indicating a good correlation, with a significance of 92% (between 2 and 3σ). Ongoing gas dispersal is certainly plausible in our clusters, given the common detection in these systems of molecular outflows (Zhang et al. 2001, 2005) which are highly effective in transferring material away from the forming objects and possibly out of the star forming region; parsec-scale flows are indeed commonly observed also from low-mass YSOs.

The final stage of gas dispersal, eventually leading to optically revealed clusters like those around Ae/Be stars, might be triggered by the powerful winds and radiation fields from newborn O and B stars; indications are (Molinari et al. 2008a, 2008b) that the most massive objects forming in the densest regions of the clumps hosting our clusters may have not yet reached the ZAMS, or are just starting to develop their HII regions. It is quite likely that this event will mark the moment of maximum efficiency of gas dispersal and further evolution of our clusters toward the Ae/Be's ones.




Fig. 16. Cluster radii (from Tab. 2) as a function of the cluster median ages (from Tab. 3). Dashed lines represent the linear fits obtained fitting in turn one plotted variable as a function of the other; the full line is the bisector of the two dashed lines and represents the fit which minimizes the quadratic geometric (i.e. not along the X or Y axis alone) distance of the data from the fit. Following Isobe et al. (1990), this is the adequate approach when the nature of the data scatter around the linear fit is not known (and it is not due to classical measurement uncertainties); the slope of the full line is -0.8 ± 0.2 and the 1σ spread is within the area enclosed by the two dashed lines.

5.2. Physical vs. Statistical Models for Cluster Formation

Figure 14 can also be used as a diagnostic to discriminate between different classes of models for the origin of clusters. Testi et al. (2001) called *physical* the class of models which imply a physical relationship between the most massive star that forms and other environmental properties like the cluster richness or the mass of the gaseous clump where the stars originate from; examples are the "turbulent core" (McKee & Tan 2003), the "coalescence" (Bonnell et al., 1998), or the competitive accretion models (Bonnell & Bate, 2006). In a second class of models, called *statistical*, the relationship between the most massive star in a cluster and its richness arises from the higher probability of finding the rare massive stars in rich clusters rather than in isolation (Bonnell & Clarke 1999). If clusters are populated by randomly picking stars from the field stars' initial mass function, and considering a cluster membership-size distribution in the form of an appropriate power law, then the observations of Testi et al. (1999) can be naturally explained. Nevertheless, this model predicts that a significant fraction of high-mass stars are still associated with relatively poorly populated clusters, in other words that massive stars can be found both in high-N clusters and, to a lesser extent, in low-N clusters.

The dashed line in Fig. 14 is the upper boundary of the region which should contain 25% of the cluster realizations obtained by randomly extracting stars from the IMF (Testi et al. 2001). Indeed, if we consider our measurements of I_c for our entire sample (i.e. the full and empty triangles), there is fraction of about 15% of the clusters which is found marginally below the dashed line. However, it has to be noted that our modeling was possible only for clusters above a membership threshold derived with I_c , it is thus a somewhat biased subsample toward rich clusters. In the extreme assumption that the fields with no detected cluster are cases of systems composed by a single heavily ex-

tinged star, and thus would fall below the dashed line in Fig. 14, then this fraction would approach the 25% level. This, however, is an extreme case because, as we already discussed, the high value of the extinction derived from submillimeter maps may be the reason for not detecting clusters in at least a number of observed fields.




Fig. 17. Mass for the highest mass star as a function of the total stellar mass for the 9 modeled clusters (see Table 3); the bars associated to each cluster (the diamond symbols) represent the total span of the parameter values for the classes of models selected by our analysis (the diamond marks the median values). The dash-dotted, and dashed lines represents the relationship obtained for numerical simulations of clusters drawn from pure *random sampling* of the IMF, and using a so-called *sorted sampling*, following Weidner & Kroupa (2006). The full line is a semi-analytical approximation of this relationship obtained by Weidner & Kroupa (2004). The dotted line is the limit where a cluster is made of just one star.

As a further diagnostic between physical and statistical cluster models, Weidner & Kroupa (2006) recently argued that a non-trivial correlation exists between the highest-mass star in a cluster, $M_{\star\text{Max}}$, and the total stellar mass of the cluster itself, $M_{\star\text{Tot}}$ (Fig. 17). Numerical simulations show that the relationship obtained by pure random sampling of the IMF with an imposed physical limit of $150 M_{\odot}$ for the maximum stellar mass (the dashed-dot line in Fig. 17) is clearly not representing our data.

A substantially different result (the dashed line) is obtained if cluster members are picked in ascending order and constrained to total cluster masses distributed according to a cluster total mass function (Weidner & Kroupa 2006). Basically, this second option represents the fact that drawing 10 cluster of 100 stars will not deliver the same $M_{\star\text{Max}}=f(M_{\star\text{Tot}})$ as drawing 1 cluster of 1000 stars. This trend closely resembles a semi-analytical approximation of $M_{\star\text{Max}}=f(M_{\star\text{Tot}})$ obtained by Weidner & Kroupa (2004), again assuming that total cluster stellar masses are distributed according to a power-law mass function. Weidner & Kroupa (2006) suggest that this *sorted sampling* way of populating a cluster can be physically understood in terms of a pre-stellar clump where initial low-amplitude perturbations start low-mass star formation; as further perturbations with larger amplitude grow, higher mass stars will start to form until the feedback from the latter will start disrupting the cloud. This scenario

of an organised star formation where low-mass stars are the first ones to form, is the same we favor (see §5.1) considering the age spread we find in our clusters in which, based on independent considerations (Molinari et al. 2008a), the most massive star may have not been formed at all. By the way, this latter possibility does not change the substance of the agreement between our data and the *physical* cluster models in Fig. 17, since the late addition of the highest mass star currently not yet visible in the near-IR would shift the points toward the top-right of the plot.

We verified *a posteriori* that the range of $M_{\star\text{Max}}$ and $M_{\star\text{Tot}}$ parameters values explored by our simulations is much wider than the area spanned by the bars attached to the single points, and also includes regions compatible with the random sampling cluster model. We then conclude that the result of Fig. 17 is not likely to be produced by a biased sampling of the clusters' physical parameters explored in our models.

The predictions of the *sorted sampling* above described, with which our data points best agree, are also in good agreement with the results from simulations of clusters forming in a competitive accretion scenario (Bonnell, Vine & Bate 2004). This model, however, seems excluded by the observed ages and age spreads in our clusters which are in clear disagreement with the predicted cluster formation timescales of $2\div3$ free-fall times.

5.3. Influence of Binarity on the Interpretation of Age Spread

Weidner, Kroupa & Maschberger (2008) carried out extensive numerical simulations to determine how the presence of unresolved binary/multiple stars can affect the observational properties of a young cluster in a massive star forming region. Assuming a 100% of binarity in a cluster of coeval sources, they find that unresolved binaries may lead an observer to conclude that instead a significant age spread is present in the cluster; the full line in Fig. 18 shows the *a posteriori* age determination assuming that all binaries are unresolved. We see that the measured age spread for the large majority of stars simulated in Weidner et al.'s simulation is comparable to a Log(age) Gaussian distribution with $\sigma=0.1$, which is one of the possible choices of Star Formation Histories in our cluster models. However, the comparison of our observed KLFs with the SCG models (Table 3) suggests age spreads much bigger than this, and more comparable to a Log(age) distribution with $\sigma=0.5$ (the dotted line in Fig. 18). We then conclude that unresolved binaries in our clusters cannot account for the observed age spread.

6. Conclusions

The main results of this work are the following:

- We have imaged in the J, H, and K_s NIR bands 26 intermediate and high-mass star forming regions selected from a larger sample of sources and spanning a range in luminosities and presumed youth. We have identified the presence of 23 young stellar clusters in 22 fields.
- Revealed clusters have richness indicator values between ten and several tens of objects and have median radius of 0.7 pc. Compared to clusters around Herbig Ae/Be stars, our clusters seem richer and larger for any given mass for the most massive star in each cluster. Color-color diagrams show that these clusters are young; many sources show colors typical of young pre-MS objects with an intrinsic IR excess arising from warm circumstellar dust. This is confirmed by the analysis of color-magnitude diagrams where a significant fraction of stars in each cluster are found in correspondence to




Fig. 18. Full line represents the age spread resulting from the simulations of Weidner, Kroupa and Maschberger (2008) for a cluster with an input age of 2×10^6 yrs and 100% binarity fraction. The two gaussians are the age weight functions used in our simulations of Gaussian Star Formation Histories, with $\sigma=0.1$ (dashed line) and 0.5 (dotted line) respectively. In case of constant SFH models, the adopted age spread is comparable to the $\sigma=0.5$ distribution in the figure.

the Pre-MS evolutionary tracks even after conservative de-reddening is applied.

- We cannot perform a direct inversion of stellar luminosities (and colors) into masses and ages; we use a Synthetic cluster Generator (SCG) model to create statistically-significant cluster simulations for different initial parameters (IMF, SFH, source ages and their distribution), and compare the synthetic KLFs and HKCFs with the observed (field star-subtracted) ones. For the fraction of clusters for which this comparison selects a well-confined region of the parameter space, we conclude that star formation in these regions *cannot* be represented with a single burst, but is a process that is spread out in time. Clusters have mean ages of a few 10^6 yrs; the ages of most of the clusters members are spread, within each cluster, between a few 10^5 yrs to a few 10^6 yrs. Together with the independent evidence that the most massive stars in these systems are very young, or not even on the ZAMS yet, this result is difficult to reconcile with any model predicting cluster formation in a crossing time.
- The cluster radii seem to be inversely proportional to their age, as also confirmed by the comparison of cluster parameters with those typical of Ae/Be systems, which are smaller and less rich. As suggested by numerical simulations in the literature, dispersal of intra-cluster gas (by, e.g., molecular outflows or radiation fields from massive stars) may lead to the loss of a fraction of cluster stellar population, thus indeed leading to smaller and less rich clusters. Our results seem in line with this prediction.
- The relation between the mass of the most massive star in a cluster and the cluster's richness indicator suggests that a *physical* rather than *statistical* nature of the cluster origin is more likely.

Acknowledgements. We thank the anonymous referee whose comments greatly helped improving the quality of the data presentation and the overall readability of the paper. We also thank P. Kroupa for his comments and suggestions.

References

- Lada, C., Adams, F. 1992, *ApJ* 393, 278
 Baumgardt, H., Kroupa, P. 2007, *MNRAS* 380, 1589
 Behrend, R. & Meader, A. 2001, *A&A* 373, 190
 Beltrán, M. T., Cesaroni, R., Neri, R., Codella, C., Furuya, R. S., Testi, L., & Olmi, L. 2004, *ApJ*, 601, L187
 Beltrán, M.T., Brand, J., Cesaroni, R., Fontani, F., Pezzuto, S., Testi, L., Molinari, S. 2006, *A&A* 447, 221
 Benjamin, R. A., Churchwell, E. B., Babler, B., et al. 2003, *PASP* 115, 953
 Bertoldi, F. & McKee, C. F. 1992, *ApJ* 395, 140
 Bessel, M. S., 1979, *PASP*, 91, 377
 Bessel, M. S. 1991, *AJ*, 102, 303
 Bessel, M. S., Brett, J.M. 1988, *PASP*, 100, 1134
 Beuther, H., Schilke, P., Sridharan, T. K., Menten, K. M., Walmsley, C. M., & Wyrowski, F. 2002, *A&A*, 383, 892
 Binney, S. J., Tremaine, S. 1987, *Galactic Dynamics*, Princeton Univ. Press, Princeton NJ
 Bonnell, I. A., Bate, M.R., & Zinnecker, H. 1998, *MNRAS*, 298, 93
 Bonnell, I. A., & Davies, M.B. 1998, *MNRAS*, 295, 691
 Bonnell, I. A., & Clarke, C. 1999, *MNRAS*, 309, 461
 Bonnell, I. A., & Bate, M.R. 2002, *MNRAS*, 336, 659
 Bonnell, I. A., & Bate, M.R. 2006, *MNRAS*, 370, 488
 Bonnell, I. A., Bate, M.R., Clarke, C., & Pringle, J.E. 2001, *MNRAS*, 323, 785
 Bonnell, I. A., Vine, S.G., & Bate, M.R. 2004, *MNRAS*, 349, 735
 Bonnell, I. A., Bate, M.R., & Vine, S.G. 2003, *MNRAS*, 343, 413
 Brand, J., Blitz, L. 1993, *A&A*, 275, 67
 Brand, J., Cesaroni, R., Palla, F., Molinari, S. 2001, *A&A*, 370, 230
 Cesaroni, R., Neri, R., Olmi, L., Testi, L., Walmsley, C. M., & Hofner, P. 2005, *A&A*, 434, 1039
 Cesaroni, R. 2005, *Ap&SS*, 295, 5
 Cesaroni, R., Galli, D., Lodato, G., Walmsley, C.M., & Zhang, Q. 2006, *Protostars & Planet V, Disks Around Young O-B (Proto)Stars: Observation and Theory*
 Chini, R., Hoffmeister, V., Kimeswenger, S., Nielbock, M., Nürnberger, D., Schmidtobreich, L., & Sterzik, M. 2004, *Nature*, 429, 155
 Clarke, C., & Delgado-Donate, E. 2003, *Galactic Star Formation Across the Stellar Mass Spectrum. ASP Conference Series*, p.287
 de Wit, W.-J., Testi, L., Palla, F., Zinnecker, H. 2005, *A&A*, 437, 247
 Hillenbrand, L. A., & Carpenter, J. M. 2000, *ApJ*, 540, 236
 Hoogerwerf, R., de Bruijne, J. H. J. & de Zeeuw, P. T. 2001, *A&A*, 365, 49
 Hartmann, L. W., Jones, B. F., Stauffer, J. R. & Kenyon, S. J. 1991, *AJ*, 101, 1050
 Hunt, L. K., Mannucci, F., Testi, L., Migliorini, S., Stanga, R. M., Baffa, C., Lisi, F., Vanzì, L., 1998, *AJ*, 115, 259
 Hillenbrand, L. A. 1997, *AJ*, 113, 1733
 Isobe, , Feigelson, E., Akritas, , Babu, , 1990, *ApJ* 364, 104
 Klein, R. I., Inutsuka, S., Padoan, P., Tomisaka, & K. 2006, *Protostars & Planet V, Current Advances in the Methodology and Computational Simulation of the Formation of Low-Mass Stars*
 Kenyon, S., & Hartmann, L. 1995, *ApJS*, 101, 117
 Keto, E. 2003, *ApJ*, 599, 1196
 Koornneef, J. 1983, *A&A*, 128, 84
 Kumar, M.S.N., Keto, E., Clerkin, E. 2006, *A&A* 449, 1033
 Kumar, M.S.N., & Grave, J.M.C. 2008, *ASPC* 387, 323
 Kroupa, P., Tout, C. A., & Gilmore, G. 1993, *MNRAS*, 262, 545
 Lada, C. J., Lada, E. A. 2003, *ARA&A* 41, 57
 McKee, C.F., Tan, J.C. 2003, *ApJ* 585, 850
 Molinari, S., Brand, J., Cesaroni, R., & Palla, F. 1996, *A&A*, 308, 573
 Molinari, S., Brand, J., Cesaroni, R., Palla, F., & Palumbo, G. G. C. 1998, *A&A*, 336, 339
 Molinari, S., Brand, J., Cesaroni, R., & Palla, F. 2000, *A&A*, 355, 617
 Molinari, S., Testi, L., Rodríguez, L.F., & Zhang, Q. 2002, *ApJ*, 570, 758
 Molinari, S., Pezzuto, S., Cesaroni, R., Brand, J., Faustini, F., Testi, L. 2008a, *A&A*, 481, 345
 Molinari, S., Faustini, F., Testi, L., Pezzuto, S., Cesaroni, R., Brand, J. 2008b, *A&A*, 487, 1119
 Moraux, E., Lawson, W. A. & Clarke, C. 2007, *A&A*, 473, 163
 Nakano, T. 1989, *ApJ*, 345, 464
 Palla, F., & Stahler, S. W. 1999, *ApJ*, 525, 722
 Preibisch, Th., Ossenkopf, V., Yorke, H. W., Henning, Th. 1993, *A&A*, 279, 577
 Rieke, G. H., & Lebofsky, M. J. 1985, *ApJ*, 288, 618
 Salpeter, E. E. 1955, *ApJ*, 121, 161
 Scalo, J. 1998, *The Stellar Initial Mass Function, ASP Conference Proceedings* Vol. 142, p. 201 (eds. G. Gilmore, D. Howell)
 Strom, K. M., Strom, S. E., Edwards, S., Cabrit, S., & Skrutskie, M. F. 1989, *AJ*, 97, 111
 Tan, J.C., Krumholz, M.R., McKee, C.F. 2006, *ApJ*
 Testi, L., Palla, F., Prusti, T., Natta, A., & Maltagliati, S. 1997, *A&A*, 320, 159

- Testi, L., Palla, F., & Natta, A., 1998, A&AS, 133, 81
- Testi, L., Palla, F., & Natta, A., 1999, A&A 342, 515
- Testi, L., Palla, F., & Natta, A., 2001, *From Darkness to Light, ASP Conference Series* Vol. 243, p. 377 (eds. T. Montmerle, Ph. André)
- Tan, J. C., 2005, *Cores to Clusters: Star Formation with Next Generation Telescopes Astrophysics and Space Science Library*, Vol. 324, p. 87
- Tan, J. C., & McKee C.F. 2002, *Hot Star Workshop III: The earliest Stages of Massive Star Birth. ASP Conference Proceedings*, 267, p. 267
- Vig. S., Testi, L., Walmsley, M., Molinari, S., Carey, S., Noriega-Crespo, A. 2007, A&A 470, 977
- Weidner, C., Kroupa, P. 2004, MNRAS 348, 187
- Weidner, C., Kroupa, P. 2006, MNRAS 365, 1333
- Weidner, C., Kroupa, P., & Maschberger, T. 2008 arXiv0811.3730W
- Yorke, H. W. 2002, *Hot Star Workshop III: The earliest Stages of Massive Star Birth. ASP Conference Proceedings*, 267, p. 165
- Zhang, Q., Hunter, T. R., Brand, J., Sridharan, T. K., Molinari, S., Kramer, M. A., Cesaroni, R. 2001, ApJ 552, L167
- Zhang, Q., Hunter, T. R., Brand, J., Sridharan, T. K., Cesaroni, R., Molinari, S., Wang, J., Kramer, M. A. 2005, ApJ 625, 864
- Zinnecker, H. & Yorke, H. W. 2007 ARA&A 45, 481

Online Material

Appendix A: K_s images for all observed fields

This appendix contains the K_s images for all observed fields; these are also available at <http://galatea.ifs-roma.inaf.it/faustini/K-images/>




Fig. A.1. K_s image of field Mol003 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.2. K_s image of field Mol008 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.3. K_s image of field Mol009; no submillimeter image is available for this field.




Fig. A.4. K_s image of field Mol011 with superimposed SCUBA 850 μm continuum in white contours (Molinari et al. 2008a).




Fig. A.5. K_s image of field Mol012 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.6. K_s image of field Mol015; no submillimeter image is available for this field.




Fig. A.7. K_s image of field Mol028; no submillimeter image is available for this field.




Fig. A.8. K_s image of field Mol030; no submillimeter image is available for this field. The white cross marks the position of the IRAS source.




Fig. A.9. K_s image of field Mol038 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.10. K_s image of field Mol045 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.11. K_s image of field Mol050; no submillimeter image is available for this field. The white cross marks the position of the IRAS source.




Fig. A.12. K_s image of field Mol059 with superimposed SCUBA 850 μm continuum in white contours (Molinari et al. 2008a).




Fig. A.13. K_s image of field Mol075 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.14. K_s image of field Mol082; no submillimeter image is available for this field. The white cross marks the position of the IRAS source.




Fig. A.15. K_s image of field Mol084; no submillimeter image is available for this field. The white cross marks the position of the IRAS source.




Fig. A.16. K_s image of field Mol098 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.17. K_s image of field Mol099; no submillimeter image is available for this field. The white cross marks the position of the IRAS source.




Fig. A.18. K_s image of field Mol103 with superimposed SIMBA 1.2mm continuum in white contours (Beltrán et al. 2006).




Fig. A.19. K_s image of field Mol109 with superimposed SIMBA 1.2mm continuum in white contours (Beltrán et al. 2006).




Fig. A.20. K_s image of field Mol110 with superimposed SIMBA 1.2mm continuum in white contours (Beltrán et al. 2006).




Fig. A.21. K_s image of field Mol136 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.22. K_s image of field Mol139 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.23. K_s image of field Mol143 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.24. K_s image of field Mol148 with superimposed MAMBO 1.1mm continuum in white contours (Molinari et al. 2008a).




Fig. A.25. K_s image of field Mol151 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).




Fig. A.26. K_s image of field Mol160 with superimposed SCUBA 850μm continuum in white contours (Molinari et al. 2008a).

Appendix B: K_s luminosity functions

This appendix presents the set of background-subtracted K_s luminosity functions for all detected clusters; the green vertical line represents the 90% completeness limit as estimated from artificial star experiments (see §2.1). Material can also be retrieved at <http://galatea.ifs-roma.inaf.it/faustini/KLF/>




Fig. B.1. Mol003 (left), Mol008A (center), Mol008B (right)




Fig. B.2. Mol009 (left), Mol011 (center), Mol012 (right)




Fig. B.3. Mol015 (left), Mol028 (center), Mol045 (right)




Fig. B.4. Mol050 (left), Mol075 (center), Mol082 (right)




Fig. B.5. Mol084 (left), Mol099 (center), Mol103 (right)




Fig. B.6. Mol136 (left), Mol110 (center), Mol136 (right)




Fig. B.7. Mol139 (left), Mol143 (center), Mol148 (right)




Fig. B.8. Mol151 (left), Mol160 (right)